Journal of Immunological Methods, 24 (1978)
© Elsevier/North-Holland Biomedical Press

SUBJECT INDEX TO VOLUME 24 (1978)

2-Aminoethylisothiouronium bromide (AET) treatment of SRBC enhances E-rosetting, 355—361
Ammonium chloride, isotonic, reduces lymphocyte cytotoxicity, 257—267
Antibody
  reaction with antigen visualized at glass surface, 57—61
  sites concentration related to affinity distribution, 63—73
  specificity in immune complex tested by radio-IE of $F(ab')_2$, 31—38
Antigen
  adsorbed to glass in visual immune reaction test, 57—61
  assayed with [^{125}I]SpA and IgG Ab bound to immobilized Ag, 269—285
  frozen tissue sections content determined by RID, 377—381
  schistosome, quantified by RIPEGA, 229—237

Bacteriophage, SEB-coated, inactivation by anti-SEB antiserum, 129—133
Blood, large volumes obtained from immunized rabbits without haematic damage, 19—24

Calcium-binding protein assayed in intestinal tissue by RID, 377—381
Chorionic gonadotropin assayed in human urine using [^{125}I]SpA-IgG Ab binding, 275—281
Complement
  activated during lymphocyte isolation from blood, 201—221
  functional C2 determined with readily available reagents, 223—227
Cytotoxicity
  ADCC and natural, reduced by NH_4Cl treatment of lymphocytes, 257—267
  allo-Ab-mediated, reveals lymphocyte chimerism in marrow-reconstituted mice, 337—343
  cell-mediated
    assayed by measuring ^{86}Rb incorporation in residual targets, 287—303
    ^{51}Cr-release microassay, 79—88
    prolonged cell incubation effects, 345—353
    in xenogeneic man-mouse response, 11—12, 14—17

Dimethylsuberimidate cross-links IgG Ab with SpA-Sepharose, 305—309

Electrophoresis, polyacrylamide gel, tests purified human α2M, 41—44
Enzyme-linked immunosorbent assay for anti-SEB antibodies, 129
Erythrocytes
  chicken and horse, generate sheep PFC in vitro, 313—319
  sheep
    AET-treated, efficiently form E-rosettes with bovine T cells, 355—361
    frozen, in HIG test for antiviral antibodies, 1—8
    sensitized, in measuring functional C2 in human sera, 224—227
  ox, anti-Ig-coupled and sheep, fluorescein-labelled, as indicator cells in IARR for Ab to T and B lymphocytes, 175—181

Farr test measurement errors statistically analyzed, 63—73

Glass
slides in visual immune reaction test, 57—61
wool filtration in purifying helper T cells, 26—28

Haemadsorption, mixed, disc test in human tissue typing, 143—144
Haemoglobin, mouse, monospecific antibody detected by IF, 383—387
Haemolysis in gel (HIG) with Ag-coupled frozen RBC detects antiviral antibodies, 1—8
Hamilton repeating syringe insures uniform droplets in LMI assay, 365—369
Hepatitis B surface (HBs) antigen, Ab subtype specificity tested by RIA, 371—375
Hypotonic lysis in purifying helper T cells, 26—27

IgG
antibodies to ragweed and grass allergens assayed by RIA using SpA, 183—191
antibody bound to immobilized antigen in assay with [^{125}I]SpA, 269—285
coupled to PVP beads via polyglutaraldehyde, 241—246
cross-linked with SpA-Sepharose using dimethylsuberimidate, 305—309
electrophoretic migration of small amounts enhanced by carrier protein, 99—110
iodoacetylated, coupled to mercaptosuccinylated ovalbumin, 321—336
IgM
assayed in human sera using [^{125}I]SpA-IgG antibody binding, 281—282
of low mol.wt. quantitated by radial immunodiffusion, 135—139
Immunoelectrophoresis (IE)
crossed, IgG migration facilitated by mixing with rabbit γ-globulin, 99—110
rocket, in the presence of denaturing agents, 155—162
Immunofluorescence (IF) detects monospecific Ab to mouse haemoglobin, 383—387
Immunosorbents, Sepharose-based
in detecting monospecific anti-haemoglobin antibody, 383—387
in isolating RNAse-free IgG, 113—114
prepared by coupling IgG to insolubilized SpA, 305—309
in purifying α2M, 40
in RIA for IgG antibodies to ragweed and grass allergens, 184—191
Indirect anti-Ig rosetting reaction (IARR) detects allo-Ab to T or non-T cells, 175—182
Influenza A antibodies detected by a modified HIG test, 2—5

Keratins examined by rocket IE in SDS, 159—161

Leukocytes
migration inhibition (LMI) assay, droplet size standardized and migration area diameters measured directly, 363—370
MN and PMN, separated by a modified one-step Hypaque-Ficoll method, 389—393
Lymphocytes
bovine T, improved E-rosetting with AET-treated SRBC, 355—361
human
alloantibodies to T or non-T cells detected by IARR, 175—182
ammonium chloride treatment reduces cytotoxicity, 257—267
Fc receptors labelled with IgG-coated PVP beads, 244—250
incubation effects on cytotoxic activity, 345—353
isolated by a modified one-step Hypaque-Ficoll method, 389—393
MLC with lymphoblastoid cells in BSA-supplemented media, 89—98
superoxide dismutase activity determined, 77—78
T, xenogeneic response to mouse splenocytes measured in vitro, 9—18

mouse
chimeric, determined by a cytotoxicity assay, 337—343
cytotoxic, studied by ^{51}Cr-release microassay, 79—88
helper T, purified and cryopreserved, 25—30
stimulation measured by [^{3}H]uridine uptake in vitro, 163—173
rabbit B and T, relative numbers altered by isolation techniques, 201—221
sheep, immune response in vitro to chicken and horse RBC, 311—320

α-2 Macroglobulin (α2M) purified by affinity chromatography, 39—45
Measles virus antibodies detected by a modified HIG test, 5—6
Methyl cellulose and carbonyl iron activate C during lymphocyte isolation, 203—220
Mixed haemadsorption (MH)
patterns obtained with selected sera compared between cell lines, 141—153
schematic procedure for disc test, 143—144
Mixed lymphocyte culture (MLC) in BSA-supplemented media, reproducible responses to blastoid cells, 89—98
Monocytes, human, incubation effects on cytotoxic activity, 345—353
Mumps virus antibodies detected by a modified HIG test, 6
Myeloma cells, mouse, proliferation measured by spleen colony-forming assay, 47—56

Opsonization activity of serum tested by electronic counting of unphagocytosed yeast, 251—256
Ovalbumin, mercaptosuccinylated, coupled to iodoacetylated IgG, 321—336

Polyglutaraldehyde couples proteins to PVP beads and labels Fc receptors, 239—250
Polymer tagging particles in visual Ag-Ab test on glass slides, 58—59
Protein-protein conjugates prepared by a new method, 321—336

Radial immunodiffusion (RID) assays antigens in frozen tissue sections, 377—381
Radial immunodiffusion quantitates low mol.wt. IgM in human sera, 135—139
Radioimmunoassay (RIA) for IgG antibodies to ragweed AgE and rye group I allergens, 183—191
Radioimmunoassay (RIA) subtypes anti-HBsAg antibodies, 371—375
Radioimmunoelectrophoresis tests Ab specificity in immune complexes, 31—38
Radioimmunoprecipitation-PEG assay RIPEGA quantifies circulating schistosome Ag, 229—237
Rosettes
E, formed efficiently by AET-treated SRBC with bovine T cells, 355—361
E, EA and EAC, in analyzing rabbit blood lymphocyte subpopulations, 201—221
Rubidium-86 incorporation in microcytotoxicity assay, 287—303

Schistosoma mansoni antigens assayed in human sera by RIPEGA, 229—237
SDS electroimmunoassay for insoluble proteins, 159—161
Spleen colonization assays idiotype-specific transplant resistance, 47—56
Staphylococcal
enterotoxin B (SEB) assayed by inhibiting antibody inactivation of SEB-bacteriophage, 127—134
protein A (SpA), insolubilized
in isolating RNAse-free IgG, 111—125
in RIA for IgG antibodies to ragweed and grass allergens, 184—191
Streptococcal protein A (SpA)
^{125}I-labelled, in assaying Ag with IgG Ab bound to immobilized Ag, 269—285
Sepharose-coupled, conjugated with IgG via dimethylsuberimidate, 305—309

Superoxide dismutase activity in human blood cells determined using logit paper, 75—78

Tissue typing of cultured human cells, MH patterns with selected sera compared in pairs, 141—153

Transplantation resistance in mice assayed by spleen colonization, 47—56

[^{3}H]Uridine uptake in vitro tests selectively stimulated lymphocytes, 163—173

X-irradiation in purifying helper T-lymphocytes, 26—28

Yeast opsonization test for immunodeficiency, 251—256